



Preliminary Report on the July 2008 Mobile Sentinel Survey in the Northern Gulf of St. Lawrence

December 2008

Sentinel Fisheries Program

Each year, biologists from Fisheries and Oceans assess the status of commercially exploited fish and invertebrate stocks. In order to perform these evaluations, scientists use data provided, in part, by mobile gear sentinel fishers (trawlers). The surveys performed by the mobile sentinel fishers cover almost the entire northern Gulf of St. Lawrence. Once analysed, the collected data are used to calculate abundance indices that are used to estimate the status of various stocks.

The mobile gear Sentinel Fisheries Program follows a depth-stratified random survey plan. The northern Gulf is divided into depth strata because depth is known to have an influence on the distribution of fish and invertebrate species. The following strata have been defined: 10-20, 20-50, 50-100, 100-150, 150-200 and over 200 fathoms (1 fathom = 1.83 meters = 6 feet). The mobile survey generally consists of 300 stations randomly selected within those strata. All strata are sampled because results from this survey are used for many species that have different depth preferences. It is also important to find the limits of distribution of a specie and to monitor any potential shifts in time. Nine trawlers, five from Newfoundland and four from Quebec perform the entire survey. At each predetermined station, the vessel performs a standard 30 minutes tow at 2,5 knots. The nine boats participating in the survey use the same trawl, a 300 Star Balloon mounted on a Rock Hopper footgear. The trawl mesh size is 145 mm with a liner of 40 mm in the codend. The use of a liner allows the sampling of fish as young as age 2 for cod but also requires the presence of an observer or a trained technician aboard each vessel. The observers are from Biorex in Quebec and Seawatch in Newfoundland. In spite of the fact that all nine boats use the same trawl, a study showed a 25% variability in wing spread opening during trawling activities. The use of a restrictor cable allows to reduce this variability from 25% to 6 % without bias due to depth and thus to compare the catches from all vessels.

The sentinel surveys require a great deal of thoroughness in collecting a variety of data. With the help of crewmembers, observers or technicians are responsible for the collection of numerous data according to a scientific protocol. At each set, the total catch is sorted by species and weighed. Then the length, weight, sex, and maturity of a number of fish of each species are recorded. For specific demands, some samples (otoliths, liver, gonads, stomach etc.) may also be taken on some fish. Moreover, fish samples are collected and frozen for diverse studies. Finally, water temperature and fishing depth data are collected using a Vemco sensor installed on the trawl. These biological and oceanographic data yield valuable information on the size, growth, condition and diet of various species, as well as stock abundance and water temperature.

The Department of Fisheries and Oceans (DFO) has primary responsibility for the administration of the sentinel program. The implementation of the program is the responsibility of the Association des Capitaines-Propriétaires de la Gaspésie inc. (ACPG) for mobile gear fisheries in Divisions 4S and 4T, and of the Fish, Food and Allied Workers (FFAW) of Newfoundland in Subdivision 3Pn and Division 4R.

July 2008 Survey

The 14th annual July sentinel survey was conducted in the northern Gulf of St. Lawrence between July 1 and July 14, 2008. A total of 289 fishing stations were successfully carried out (Figure 1), i.e. 21 in 3Pn, 129 in 4R (including 10 tows in the 10 to 20 fathoms strata), 112 in 4S and 27 in 4T. Those 289 stations represent 96% of the sampling target.

From July 2nd to July 14th, four Quebec trawlers sampling 4ST completed 139 out of a planned 152 stations (Figure 1). On the west coast of Newfoundland (3Pn, 4R), five trawlers carried out all 150 planned stations from July 1st to July 4th (Figure 1).

The 27 tows done in the 4T Division are conducted to complement the assessment of the Unit 1 redfish, Greenland halibut (turbot) and Atlantic halibut stocks of the management unit 4RST. **The cod catches in 4T are not used to estimate the abundance of the northern Gulf of St-Lawrence cod stock (3Pn, 4RS).**

Catches from the 10 tows carried out in the three coastal strata in 4R (10 - 20 fathoms) were used to calculate an index of minimum trawlable biomass for cod. However, the treatment of this new index (2003 to 2008) will be re-evaluated in the upcoming cod assessment in February 2009.

Cod, redfish, turbot and Atlantic halibut catches for the 289 successful tows of the 2008 July survey are presented in table 1. This table also shows the total catches including the unsuccessful and the discretionary tows. It is important to note that following the Larocque case, no discretionary tows have been allowed since 2006. In 2008, the fish composing the main proportion of the other fish species category are American plaice, herring and thorny skate.




Figure 1: Distribution of stratified random tows performed during the July 2008 survey.

Table 1: Cod, redfish, turbot and Atlantic halibut catches for the successful tows and total catches including unsuccessful and discretionary tows for the July 1995-2008 surveys (3Pn, 4RST).

| Year | Sets Survey | Survey Catches (kg) | | | | Sets Total | Total Catches (kg) | | | | |
|-------|----------------|---------------------|---------|--------|---------|---------------|--------------------|---------|--------|---------|---------------|
| | | Cod | Redfish | Turbot | Halibut | | Cod | Redfish | Turbot | Halibut | Other Species |
| 1995 | 311 | 6 477 | 11 457 | 649 | 84 | 326 | 6 598 | 11 662 | 675 | 84 | 4 716 |
| 1996 | 272 | 7 254 | 16 921 | 1 300 | 114 | 332 | 12 108 | 27 169 | 1 502 | 150 | 8 593 |
| 1997 | 285 | 8 642 | 12 358 | 1 206 | 27 | 313 | 11 271 | 13 582 | 1 397 | 80 | 5 848 |
| 1998 | 289 | 7 719 | 16 154 | 1 472 | 17 | 320 | 12 196 | 36 231 | 1 668 | 113 | 7 198 |
| 1999 | 294 | 5 487 | 12 623 | 1 703 | 42 | 335 | 19 396 | 17 177 | 2 079 | 129 | 4 031 |
| 2000 | 291 | 7 893 | 7 574 | 1 583 | 97 | 324 | 16 963 | 10 486 | 1 932 | 126 | 5 454 |
| 2001 | 275 | 10 238 | 7 603 | 1 342 | 120 | 317 | 16 476 | 14 421 | 1 814 | 208 | 4 194 |
| 2002 | 261 | 7 729 | 8 101 | 1 486 | 113 | 293 | 18 551 | 8 849 | 3 090 | 160 | 4 155 |
| 2003 | 296 | 13 741 | 6 400 | 1 693 | 44 | 326 | 14 040 | 6 616 | 3 512 | 72 | 3 590 |
| 2004 | 280 | 14 072 | 8 245 | 2 015 | 216 | 317 | 15 655 | 13 295 | 2 567 | 271 | 6 670 |
| 2005 | 285 | 9 662 | 6 785 | 2 977 | 226 | 303 | 10 023 | 7 802 | 3 649 | 402 | 8 652 |
| 2006 | 295 | 13 174 | 5 106 | 2 748 | 335 | 325 | 15 332 | 5 963 | 3 624 | 577 | 6 647 |
| 2007* | 291 | 6 431 | 6 797 | 2 976 | 382 | 297 | 6 435 | 6 836 | 2 977 | 399 | 3 905 |
| 2008* | 289 | 9 931 | 4 310 | 2 594 | 456 | 293 | 9 931 | 4 341 | 2 604 | 456 | 2743 |

* No discretionary tows

1. Biomass and Distribution of Groundfish

Cod

The July sentinel survey series (1995-2008) suggests an increase in the minimum trawlable biomass for cod between 1995 and 2000 with a rather stable period up to 2006. Data for 2007 and 2008 indicate a marked decrease in the biomass index which is comparable to the level observed in 1995 (Figure 2a). The biomass estimate for 2008 in Subdivision 3Pn and Divisions 4RS (more than 20 fathoms) is 33 235 tons and it is the second lowest value since the beginning of the sentinel mobile survey.




Figure 2: Minimum trawlable biomass index for cod based on the stratified random tows of the July sentinel mobile survey in Subdivision 3Pn and Divisions 4RS (1995 – 2008). a) For strata of depth greater or equal to 20 fathoms b) Including the three strata in 10-20 fathoms.

In 2008, for waters deeper than 20 fathoms, the minimum trawlable biomass for 4S shows a decrease of 63% compared to 2007, going from 11 468 tons to 4 264 tons. This latest value is similar to those observed from 2000 to 2006. For the 4R Division, the biomass estimate for 2008 is 28 931 tons, which is very similar to the 2007 estimate and much lower than the 2000 to 2006 values. The 3Pn estimate in 2008 is the lowest of the 1995-2008 series with 40 tons.

Since 2003, three inshore strata with depth range of 10 to 20 fathoms are sampled in the 4R Division. The purpose of adding those strata was to look for the presence of cod outside the zone previously sampled by trawlers in the July mobile gear sentinel survey. The location of the strata are as follows: one on both side of the strait of Belle Isle; one north of the 49th parallel; and a third one south of the 49th parallel. Ten tows were done in these strata by five trawlers during the July 2008 survey (Figure 3). The cod catches varied between 0 and 1 566 kg for a 30-minutes standard tow. Nine of the ten tows reached the 30 minutes duration while the other one lasted less than 30 minutes because of bad bottom (trawl hooked at the bottom). This year, the minimum trawlable biomass for those three strata is the highest of the short series with 34 391 tons (Figure 2b). The extension of the survey area to sample waters of 10 to 20 fathoms was thus justified and will be maintained in the future. The 2008 estimate of total minimum trawlable biomass (including the 10 to 20 fathoms strata) is 67 627 tons, which is slightly higher than the 2007 estimate but still lower than the other values in the short series.

As in the past, the catch distribution shows that cod is located primarily in 4R Division along the west coast of Newfoundland (Figure 3). In 2008, the cod concentration remains low in 4S Division and 3Pn Subdivision. Of the 289 tows performed in this survey, 7 had catches of cod of more than 500 kg and they were all located in the 4R Division. In addition, 5 of those tows were done in the strata with depth range between 10 and 20 fathoms. The largest catches for 4S and 3Pn are 78 and 3.5 kg respectively.




Figure 3: Bottom temperature and observed catch rate (kg / standard tow) distribution of cod for the July 2008 stratified random survey in 3Pn Subdivision and 4RST Divisions. Catches greater than 500 kg are identified on the map.

Redfish

For Unit 1 redfish stock (4RST Divisions), the July sentinel survey series (1995-2008) indicates a higher minimum biomass estimate between 1996 and 1999. In 2000, there is a decrease in the minimum trawlable biomass index which is relatively stable up to 2008 (figure 4).

As in earlier years, the redfish was concentrated for the most part in the channels of the Northern region of the Gulf (Figure 5). In July, good concentrations of redfish were found in the 3Pn Subdivision. This area is not part of the redfish stock of Unit 1 management (4RST) between the months of June to December.




Figure 4: Minimum trawlable biomass index for redfish in 4RST based on the July stratified random survey (1995 – 2008).




Figure 5: Bottom temperature and catch rate (kg / standard tow) distribution of redfish for the July 2008 stratified random survey in 3Pn and 4RST.

Turbot (Greenland Halibut)

For the Gulf of St. Lawrence turbot stock (4RST Divisions), the July sentinel survey series (1995 – 2008) shows a general increase in biomass from 1995 to 2005 with a relative stability up to 2008 (Figure 6). The total biomass estimate for 2008 is the third highest of the series with 34 561 tons.

Turbot was concentrated mostly in the Estuary and in the Laurentian Channel, around Anticosti Island and in the Northern portion of the Esquiman Channel (Figure 7). The distribution of Greenland halibut is overall similar to those of earlier years. The mobile sentinel survey does not sample the Estuary where turbot is found in abundance in the August DFO's annual scientific survey and in the commercial fishery.




Figure 6: Minimum trawlable biomass index for Greenland halibut based on the July stratified random survey (1995–2008).




Figure 7: Bottom temperature and catch rate (kg/standard tow) distribution of Greenland halibut for the July 2008 stratified random survey in area 3Pn, 4RST.

Atlantic Halibut

The minimum trawlable index for Atlantic halibut based on the sentinel survey shows a low and stable biomass between 1995 and 2003 with a gradual increase up to 2007. The biomass estimate slightly decreases to 3 345 tons in 2008, which is similar to the 2006 estimate (Figure 8).

The catch distribution map of Atlantic halibut shows its presence in the Esquiman, Laurentian and Anticosti channels (Figure 9).




Figure 8: Minimum trawlable biomass index for Atlantic halibut based on the July stratified random survey (1995–2008).




Figure 9: Bottom temperature and catch rate (kg/standard tow) distribution of Atlantic halibut for the July 2008 stratified random survey in area 3Pn, 4RST.

2. Sampling and Analysis

The catches were sorted by species, weighted and/or counted. The sampling of biological data (length, sex and weight) was completed for cod, redfish, Greenland halibut, Atlantic halibut and witch flounder.

Otoliths

Otoliths were taken from cod in area 3Pn, 4RS. The otoliths are used to determine the age of individual specimens and these data will be included in the next cod stock assessment to be held in winter of 2009. Age is one of the key parameter in the study of fish population dynamics and in the evaluation of the abundance for the northern cod of St. Lawrence (3Pn, 4RS). The abundance is actually assessed with a model that has age-structured data as inputs. Such models allow a better monitoring of the evolution of the stock dynamic. These models are named Sequential Population Analysis models (SPA).

Stomachs

Cod and Atlantic halibut stomachs were collected to study the fish diet. Frozen samples were sent to the Maurice Lamontagne Institute where Denis Chabot's group will analyse the stomach contents.

Witch Flounder

In addition to the total weight per set, length frequencies per sex were collected for the witch flounder's assessment. Douglas Swain of the Gulf Fisheries Center in Moncton (DFO) is the scientist responsible for the stock assessment of that specie.

Acknowledgements

We wish to acknowledge the dedicated work of all the skippers, crewmembers, observers, technicians, and coordinators who contributed in reaching the objectives of the 14th annual July sentinel survey.

Table 2. List of the skippers (*boat name*), crewmembers, technicians, and observers (company) who contributed to the July 2008 sentinel survey:

| 3Pn, 4R | | | 4S, 4T | | |
|---|---|---|---|---|------------------------------|
| Skipper | Crew | Observer | Skipper | Crew | Observer |
| Winsor Hedderson (<i>Northern Tip</i>) | Derrick Pittman Anthony Pilgram Jason Saunders | Wade Saunders (FFAW) Angus Fillier (Seawatch) | Jean-Pierre Élément (<i>Rémy Martin</i>) | Rémy Élément Martin Élément | André Gagnon (Biorex) |
| Garfield Warren (<i>885-77</i>) | Leonard Warren Enos Gaulton Ephrim Smith | Monty Way (FFAW) | Clément Samuel (<i>Sciène</i>) | Normand Samuel Johnny Dumaresq | André Rioux (Biorex) |
| Dereck Coles (<i>Catalina Venture</i>) | Bob Campbell Ashley Coles Gorvin Williams Abe Coles Floyd Genge | Jason Spingle (FFAW) | Marcel Roy (<i>Sextan</i>) | Gildas Cotton Jean-Guy Côté Mathieu Roy | Christian Girard (Biorex) |
| Murray Lavers (<i>Sylvia Lynn II</i>) | Floyd Biggin Warren House Barry Ryan Thomas Lavers | Brent Hedderson (FFAW) Mike Colbourne (Seawatch) | Réjean Bernatchez (<i>Chlorydon</i>) | Paul-René Côté Gilles Côté | Marcel Boucher (Biorex) |
| Dan Genge Jr. (<i>NFLD Storm</i>) | Kevin Genge Hank Poole Genna Genge Jamie Genge | Levi Harvey (Seawatch) | | | |

For more information:

Bérubé, M. and A. Fréchet. 2001. Summary of the northern Gulf sentinel tagging program with emphasis on recaptures from adjacent management units. CSAS, Res. Doc. 2001/002. 24p.

http://www.dfompo.gc.ca/csas/csas/DocREC/2001/RES2001_002b.pdf

Bourdages, H., D. Archambault, B. Bernier, A. Fréchet, J. Gauthier, F. Grégoire, J. Lambert and L. Savard. 2008. Preliminary results from the groundfish and shrimp multidisciplinary survey from August 2007 in the northern Gulf of St. Lawrence. CSAS, Res. Doc. 2008/046. 77p

http://www.dfo-mpo.gc.ca/CSAS/Csas/Publications/ResDocs-DocRech/2008/2008_046_e.pdf

Campana, S., G. Chouinard, M. Hanson and A. Fréchet. 1999. Mixing and migration of overwintering Atlantic cod stocks near the mouth of the Gulf of St. Lawrence. Can. J. Fish. Aquat. Sci. 56 : 1873-1881.

DFO, 2008. Assessment of cod stock in the Northern Gulf of St-Lawrence (3Pn, 4RS) in 2007. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2008/003.

http://www.dfo-mpo.gc.ca/CSAS/Csas/Publications/SAR-AS/2008/SAR-AS2008_003_E.pdf

Dutil, J.-D., J. Gauthier, A. Fréchet, M. Castonguay, Y. Lambert, and P. Ouellet. ICES. 2005. Spawning and life history information for North Atlantic cod stocks. Ices Cooperative Research Report, No. 274.152 p

Dutil, J.-D., J. Gauthier, Y. Lambert, A. Fréchet and D. Chabot. 2003. Cod stocks rebuilding and fish bioenergetics : low productivity hypothesis. CSAS, Res. Doc. 2003/060. 43p.

http://www.dfo-mpo.gc.ca/csas/Csas/DocREC/2003/RES2003_060_E.pdf

Fréchet, A. 2005. Changes in biological characteristics of the fisheries of the Gulf of St. Lawrence (NAFO Divisions 4RST) since 1960. CSAS, Res. Doc. 2005/088. 34p.

http://www.dfo-mpo.gc.ca/csas/csas/publications/resdocs-docrech/2005/2005_088_e.htm

Fréchet, A., J. Gauthier, P. Schwab, H. Bourdage, C. Tournois, J. Spingle, M. Way and F. Collier. 2007. The status of cod in the Northern Gulf of St. Lawrence (3Pn, 4RS) in 2006. CSAS, Res. Doc. 2007/068. 89p.

http://www.dfo-mpo.gc.ca/csas/Csas/Publications/ResDocs-DocRech/2007/2007_068_e.htm

Fréchet, A., R. Butler, E. Hussey, J. Kennedy and G. Ross. 2003. Quality at landing in relation to gears used in the 3Pn, 4R cod fishery, 2000-2002. CSAS, Res. Doc. 2003/066. 13p.

http://www.dfo-mpo.gc.ca/csas/csas/DocREC/2003/RES2003_066_B.pdf

Fréchet, A., 1997. Standardization of otter trawlers participating in the sentinel fisheries in the northern Gulf of St. Lawrence. DFO, CSAS, Res. Doc. 97/72. 10 p.

Fréchet, A., 1996. Intercalibration of eight otter-trawlers participating in the sentinel fisheries in the northern Gulf of St. Lawrence (3Pn, 4RS) in 1995 through the use of SCANMAR sensors. DFO, Atlantic Fisheries, Res. Doc. 96/67. 15 p.

Gillis, D. J. 2002. Workshop on the Groundfish Sentinel Program. Moncton, New Brunswick, November 7- 9 2001. CSAS, Proceedings Series 2002/003. 93p.

http://www.dfo-mpo.gc.ca/csas/Csas/Proceedings/2002/PRO2002_003e.pdf

Grégoire, F., and A. Fréchet. 2005. Calculation of Northern Gulf of St. Lawrence cod (*Gadus morhua*) natural mortality for the 1990 to 2004 period. CSAS, Res. Doc. 2005/019. 25p.

http://www.dfo-mpo.gc.ca/csas/Csas/DocREC/2005/RES2005_019_b.pdf

Hammill, M.O. and G.B. Stenson. 2000. Estimated prey consumption by harp seals (*Phoca groenlandica*), grey seals (*Halichoerus grypus*), harbour seals (*Phoca vitulina*) and hooded seals (*Cystophora cristata*). J. Northw. Atl. Fish. Sci. 26: 1-23, 2000.

Rice, J.C., P.A. Shelton, D. Rivard, G.A. Chouinard and A. Fréchet. 2003. Recovering Canadian Atlantic Cod Stocks : The Shape of Things to Come?. ICES C.M., 2003/U:06.

Savenkoff, C., M. Castonguay, D. Chabot, A. Fréchet, M.O. Hammill, and L. Morissette. 2006. Main prey and predators and estimates of mortality of Atlantic cod (*Gadus morhua*) in the northern Gulf of St. Lawrence during the mid-1980s, mid-1990s, and early 2000s. Can. Tech. Rep. Fish. Aquat. Sci. 2666: vi+ 32 pp.

Shelton, P.A., J.C. Rice, D. Rivard, G.A. Chouinard and A. Fréchet. 2003. Recent progress on the implementation of the precautionary approach on Canadian cod stocks leading to the re-introduction of the moratorium. ICES C.M., 2003/Y:15.

Smedbol, R.K., P.A. Shelton, D.P. Swain, A. Fréchet and G.A. Chouinard. 2002. Review of population structure, distribution and abundance of cod (*Gadus morhua*) in Atlantic Canada in a species-at-risk context. CSAS. Res. Doc. 2002/082. 134p.

http://www.dfo-mpo.gc.ca/csas/csas/DocREC/2002/RES2002_082e.pdf

Stenson, G.B. and M.O. Hammill. 2004. Quantifying uncertainty in estimates of Atlantic cod (*Gadus morhua*) consumption by harp seals (*Phoca groenlandica*). Secr. can. consult. sci. du MPO. Doc. Rech. 2004/089 p.32.

Yvelin, J.-F., A. Fréchet, and J.-C. Brêthes. Migratory routes and stock structure of cod from the Northern Gulf of St. Lawrence (3Pn, 4RS). CSAS. Res. Doc. 2005/055. 50 p.

http://www.dfo-mpo.gc.ca/csas/Csas/DocREC/2005/RES2005_055_e.pdf

For more information, you can contact:

This report is available at our
web site

DFO

<http://www.osl.gc.ca>

Alain Fréchet,
Cod Biologist (418) 775-0628

Johanne Gauthier,
Sentinel fisheries Biologist (418) 775-0871

Philippe Schwab,
Cod Technician (418) 775-0626

ACPG inc.

Jean-Pierre Couillard
Executive Director (418) 269-7701

Coralie Tournois,
Scientific coordinator (418) 724-4658

F.F.A.W.

David Decker,
Secretary treasurer (709) 576-7276

Jason Spingle,
Staff representative (709) 634-7382

Monty Way
Scientific coordinator (709) 634-7382

Myra Swyers,
Data coordinator (709) 634-7382

SENTINEL FISHERIES PROGRAMS IN
THE NORTHERN GULF OF ST. LAWRENCE

A partnership
between
fishermen
and
Fisheries and Oceans Canada

Canada

FFAW FISH, FOOD AND ALLIED WORKERS

CAW CANADIAN ALLIED WORKERS

www.osl.gc.ca

Association des Capitaines-Propriétaires de la Gaspésie ACPG

Fisheries and Oceans Pêches et Océans Canada